



NeuroDose 1P-LSD

Product Information Guide

This guide is designed to ensure the safety of those who decide to use this substance legally.

Active ingredients: 1P-LSD (1-Propionyl-Lysergic Acid Diethylamide) 1ug (micrograms) per 1ml (millilitres) distilled water.

Mixing Directions: Place the 1P-LSD tab into the bottle of medical grade distilled water. DO NOT touch the tab with your fingers AT ALL. You can put it directly into the bottle from the plastic bag or use tweezers.

Shake well and refrigerate at least 24 hours before use. Shake well before each use.

Uses: 1P-LSD may help with depression, cerebral blood flow (CBF), smoking cessation, anxiety, and eating disorders as well as increase flow state, libido, creativity, compassion, empathy, social interactions and may boost confidence & self esteem

Warnings

Do not take this product if you are currently taking prescription medications such as Tricyclic antidepressants, serotonin reuptake inhibitors (SSRIs), serotonin-norepinephrine reuptake inhibitors (SNRIs) or lithium bicarbonate as these could cause a dangerous condition known as serotonin syndrome. It is not advised to use this product if you are taking monoamine oxidase inhibitors (MAOIs) or antipsychotics can diminish the effects of psychedelics in general.

Do not simultaneously use these types of medicines in combination with the provided 1P-LSD in this microdosing kit or any other psychedelic. Mixing 1P-LSD with alcohol or any other recreational drugs is unsafe and is not recommended.

If you exceed the maximum dose of 1P-LSD, being 15ug/ml you should not drive or operate heavy machinery for at least 12 hours after ingestion.

Side effects reported using 1P-LSD at Dosage Levels Ranging from 20ug/ml - 100ug/ml

When using 1P-LSD some of the possible side effects may include: disorganized thinking, increased emotional sensitivity, increased anxiety, increased energy, euphoria, restlessness, increased creativity, visual hallucinations, and impaired judgement

Drug Class: 1P-LSD is a hallucinogenic drug, which means that it causes subjective changes to consciousness, emotions, and thoughts.

Directions: The suggested dose of 1P-LSD is 7-12ml or 7-12 micrograms. Begin with 1-5ml/ug as a "reactant dose" to see if you have any side effects or allergies.

Other Information: To avoid building a tolerance to 1P-LSD, **we suggest microdosing every 4th day/morning** i.e if you microdose Monday morning, do not microdose again until Thursday morning, or every 96 hours. The best time to microdose is first thing in the morning.

1P-LSD can take anywhere from 30-90 minutes to take effect and stay active in the body for 8-12 hours.

Inactive Ingredients: Distilled Water

Disclaimer: While many people who microdose 1P-LSD and 4-ACO-DMT have reported positive improvements to mood, addictions, depression, cluster headaches, cerebral blood flow (CBF), smoking cessation, libido, anxiety disorders, flow state, creativity, compassion, empathy, social interactions, confidence and self esteem. NeuroDose makes no claims towards any efficacy of these products to treat any medical conditions or mental health disorders.

This product is not sold for human consumption or veterinarian use.